

# VERTEX REPRESENTATIONS FOR TWISTED AFFINE LIE ALGEBRA OF TYPE $A_{2l}^{(2)}$

LI-MENG XIA\*, NAIHONG HU\*, AND XIAOTANG BAI

**ABSTRACT.** In this paper, we construct an irreducible vertex module for twisted affine Lie algebra of type  $A_{2l}^{(2)}$ .

*Key Words:* Vertex representation, twisted affine Lie algebra,  $q$ -character

## 0. INTRODUCTION

Since the first vertex construction was discovered by Lepowsky and Wilson (1978), the vertex representations for any (untwisted) affine Lie algebra have been constructed on certain Fock space by many authors. Particularly, the vertex representations for non-simply-laced cases in [XH] are given through the twisted Heisenberg algebras. In fact, those modules are irreducible for certain twisted affine Lie algebras. Consequently, the irreducible vertex modules have been given for  $A_{2l-1}^{(2)}, D_l^{(2)}, E_6^{(2)}, D_4^{(3)}$ . However, the vertex representations for  $A_{2l}^{(2)}$  are not known yet.

In this paper, we will give an explicit construction for  $A_{2l}^{(2)}$  through its Heisenberg subalgebra. Moreover, such vertex modules are also irreducible.

## 1. TWISTED AFFINE LIE ALGEBRA OF TYPE $A_{2l}^{(2)}$

Suppose that  $\mathcal{G}$  is a complex simple Lie algebra of type  $A_{2l}$  and  $\sigma$  its diagram automorphism with order 2. Let

$$\mathcal{G}_i = \{x \in \mathcal{G} \mid \sigma(x) = (-1)^i x\}$$

for  $i \in \mathbb{Z}$ . Then

$$\mathcal{G} = \mathcal{G}_0 \oplus \mathcal{G}_1$$

as  $\mathbb{C}$ -spaces and  $\mathcal{G}_0$  is a simple Lie algebra of type  $B_l$ ,  $\mathcal{G}_1$  is an irreducible  $\mathcal{G}_0$ -module isomorphic to  $L(\Lambda_1)$ .

Let  $s$  be an indeterminate, then the linear space

$$\mathcal{G}^\sigma = \sum_{i \in \mathbb{Z}} \mathcal{G}_i \otimes s^{\frac{i}{2}} \oplus \mathbb{C}c \oplus \mathbb{C}d$$

---

\*L.M., supported by the Science Foundation of Jiangsu University (Grant No.07JDG035).

\*N.H., Corresponding Author, supported in part by the NNSF (Grants 10431040, 10728102), the PCSIRT, the National/Shanghai Priority Academic Discipline Programmes.

is an affine Lie algebra of type  $A_{2l}^{(2)}$  with Lie bracket

$$\begin{aligned} (1) \quad & [x \otimes s^m, y \otimes s^n] = [x, y] \otimes s^{m+n} + \delta_{m+n,0} m(x, y)c, \\ (2) \quad & [d, x \otimes s^m] = mx \otimes s^m, \\ (3) \quad & [c, \mathcal{G}^\sigma] = 0. \end{aligned}$$

Here  $m, n \in \frac{1}{2}\mathbb{Z}$  and  $(\cdot, \cdot)$  is a non-degenerate invariant bilinear form on  $\mathcal{G}$ .

Let  $\mathcal{H}_0$  be a Cartan subalgebra of  $\mathcal{G}_0$  and  $\alpha_1, \dots, \alpha_l \in \mathcal{H}_0^*$  be such that

$$(4) \quad (\alpha_i, \alpha_i) = \begin{cases} \frac{1}{2}, & i = l, \\ 1, & i < l, \end{cases}$$

and

$$(5) \quad (\alpha_i, \alpha_j) = \begin{cases} -\frac{1}{2}, & |i - j| = l, \\ 0, & |i - j| > l, \end{cases}$$

so

$$\Pi = \{\alpha_i \mid i = 1, \dots, l\}$$

is an prime root system of  $\mathcal{G}_0$ . Let  $\mathcal{Q} = \text{Span}_{\mathbb{Z}}\{\alpha_1, \dots, \alpha_l\}$ . Note that there is a linear isomorphism

$$\begin{aligned} \gamma : \mathcal{H}_0 &\longrightarrow \mathcal{H}_0^* \\ \alpha_i^\vee &\longmapsto \frac{2\alpha_i}{(\alpha_i, \alpha_i)}. \end{aligned}$$

Obviously,  $\mathcal{H}^\sigma = \mathcal{H}_0 \oplus \mathbb{C}c \oplus \mathbb{C}d$  is a Cartan subalgebra of  $\mathcal{G}^\sigma$ . Extend the bilinear form of  $\mathcal{H}_0$  to  $\mathcal{H}^\sigma$  via

$$(6) \quad (c, d) = 1, \quad (c, \alpha_i) = (d, \alpha_i) = 0, \quad i = 1, \dots, l.$$

Set  $\beta = \frac{(c+d)}{\sqrt{2}} + \alpha_l$ , then  $(\beta, \beta) = \frac{3}{2}$  and  $(\beta, \alpha_i) = (\delta_{i,l} - \delta_{i,l-1})\frac{1}{2}$ .

**Lemma 1.1.** *Suppose that  $\dot{\Delta}$  is the root system of  $\mathcal{G}_0$  and  $\dot{\Delta}_S$  the subset of all short roots. Then  $\mathcal{G}^\sigma$  has a root system*

$$\Delta = \{n\delta \pm \alpha, (2n+1)\delta \pm 2\alpha', m\delta \mid n \in \mathbb{Z}, m \in \mathbb{Z} \setminus \{0\}, \alpha \in \dot{\Delta}, \alpha' \in \dot{\Delta}_S\}.$$

For convenience, define

$$\begin{aligned} \Delta_L &= \{2\alpha \mid \alpha \in \dot{\Delta}_S\}, \\ \Delta_M &= \{\alpha \mid \alpha \in \dot{\Delta} \setminus \dot{\Delta}_S\}, \\ \Delta_S &= \{\alpha \mid \alpha \in \dot{\Delta}_S\}. \end{aligned}$$

Denoted by  $\mathcal{Q}_M$  the lattice generated by  $\alpha_i$  ( $0 < i < l$ ).

## 2. VERTEX MODULE

Let  $H = \mathbb{C} \otimes \mathcal{Q}$ ,  $H_M = \mathbb{C} \otimes (\mathcal{Q}_M + \mathbb{Z}\beta)$  and  $H(n), H_M(n + \frac{1}{2})$  be their isomorphic copies for  $n \in \mathbb{Z}$ , respectively. Then

$$\tilde{H} = \bigoplus_{n \in \mathbb{Z}} H(n) \oplus \bigoplus_{n \in \mathbb{Z}} H_M\left(n + \frac{1}{2}\right) \oplus \mathbb{C}c$$

is a Lie algebra with brack

$$(7) \quad \left[ \alpha' \left( m + \frac{1}{2} \right), \alpha'' \left( n - \frac{1}{2} \right) \right] = \delta_{m+n,0} \left( m + \frac{1}{2} \right) (\alpha', \alpha'') c,$$

$$(8) \quad [a'(m), a''(n)] = \delta_{m+n,0} m (\alpha', \alpha'') c,$$

$$(9) \quad [\tilde{H}, c] = 0,$$

where  $\alpha', \alpha'' \in H_M, a', a'' \in H$ , and it has a Heisenberg subalgebra

$$\hat{H} = \bigoplus_{n \in \mathbb{Z} \setminus \{0\}} H(n) \oplus \bigoplus_{n \in \mathbb{Z}} H_M \left( n + \frac{1}{2} \right) \oplus \mathbb{C}c,$$

and an abelian subalgebra

$$\widehat{H}^- = \bigoplus_{n \in \mathbb{Z}^-} H(n) \oplus \bigoplus_{n \in \mathbb{Z}^-} H_M \left( n + \frac{1}{2} \right).$$

Let  $\mathbb{C}[\mathcal{Q}]$  be the space linearly generated by  $e^{\alpha+\lambda} (\alpha \in \mathcal{Q})$  and  $S(\widehat{H}^-)$  be the symmetric algebra generated by  $\widehat{H}^-$ . Where

$$\lambda = \sum_{i=1}^l \frac{i}{2} \alpha_i.$$

Then we have

$$(\lambda, \alpha_i) = \frac{1}{4} \delta_{i,l}.$$

Define

$$V(\mathcal{Q}) = S(\widehat{H}^-) \otimes \mathbb{C}[\mathcal{Q}].$$

**Theorem 2.1.**  $V(\mathcal{Q})$  is a  $\tilde{H}$ -module defined by

$$(10) \quad a \left( -\frac{n}{2} \right) \cdot (v \otimes e^{\alpha+\lambda}) = a \left( -\frac{n}{2} \right) v \otimes e^{\alpha+\lambda}, n \in \mathbb{Z}^+,$$

$$(11) \quad b(-n) \cdot (v \otimes e^{\alpha+\lambda}) = b(-n) v \otimes e^{\alpha+\lambda}, n \in \mathbb{Z}^+,$$

$$(12) \quad b(0) \cdot (v \otimes e^{\alpha+\lambda}) = (b, \alpha + \lambda) v \otimes e^{\alpha+\lambda},$$

$$(13) \quad c \cdot (v \otimes e^{\alpha+\lambda}) = v \otimes e^{\alpha+\lambda},$$

and  $a(n - \frac{1}{2}), b(n)(n > 0)$  act as partial differential operators for which

$$(14) \quad a \left( n - \frac{1}{2} \right) \cdot a' \left( m + \frac{1}{2} \right) = \delta_{m+n,0} \left( n - \frac{1}{2} \right) (a, a'),$$

$$(15) \quad b(n) \cdot b'(m) = n \delta_{m+n,0} (a, a'),$$

for  $a, a' \in H_M, b, b' \in H$  and  $\alpha \in \mathcal{Q}$ .

### 3. 2-COCYCLE

Define bilinear map  $\epsilon : \mathcal{Q} \longrightarrow \{\pm 1\}$  by

$$(16) \quad \epsilon \left( \sum_{i=1}^l k_i \alpha_i, \sum_{j=1}^l r_j \alpha_j \right) = \prod_{i,j=1}^l \epsilon(\alpha_i, \alpha_j)^{k_i r_j},$$

$$(17) \quad \epsilon(\alpha_i, \alpha_j) = \begin{cases} -1, & i = j + 1 \\ 1, & \text{otherwise.} \end{cases}$$

**Lemma 3.1.** *If  $\alpha, \alpha', \alpha + \alpha' \in \dot{\Delta}$ , then*

$$(18) \quad \epsilon(\alpha, \alpha') = -\epsilon(\alpha', \alpha).$$

*Proof.* For the definition of this map, one can see [1], [4]. In fact it is a (sign of structure constants) 2-cocycle of  $B_l$  type.  $\square$

Define a map  $p : \mathcal{Q} \longrightarrow H_M$  by

$$(19) \quad p \left( \sum_{i=1}^l k_i \alpha_i \right) = \sum_{i=1}^{l-1} \text{sgn}(k_i) \left( k_i - 2 \left\lfloor \frac{k_i}{2} \right\rfloor \right) \alpha_i + \text{sgn}(k_l) \left( k_l - 2 \left\lfloor \frac{k_l}{2} \right\rfloor \right) \beta.$$

where  $\text{sgn}(k) = 1$  if  $k \geq 0$  and  $\text{sgn}(k) = -1$  if  $k < 0$ . Also define

$$(20) \quad p_0 \left( \sum_{i=1}^l k_i \alpha_i \right) = \sum_{i=1}^l \text{sgn}(k_i) \left( k_i - 2 \left\lfloor \frac{k_i}{2} \right\rfloor \right) \alpha_i.$$

**Lemma 3.2.** (1) *If  $\alpha \in \Delta_L$ , then  $p(\alpha) = p_0(\alpha) = 0$ ;*

(2) *If  $\alpha \in \Delta_M$ , then  $p(\alpha) = p_0(\alpha) = \pm(\alpha_i + \alpha_{i+1} + \cdots + \alpha_j)$  for some  $1 \leq i < j < l$ ;*

(3) *If  $\alpha \in \Delta_S$ , then*

$$p(\alpha) = \pm(\alpha_i + \alpha_{i+1} + \cdots + \alpha_{l-1} + \beta)$$

*for some  $1 \leq i \leq l$ , and  $p_0(\alpha) = \alpha$ .*

#### 4. VERTEX CONSTRUCTION

Let  $z$  be a complex variable. For  $\alpha, r \in \mathcal{Q}$ , define  $\mathbb{C}$ -linear operators as

$$z^\alpha(v \otimes e^{r+\lambda}) = z^{(\alpha, r+\lambda)} v \otimes e^{r+\lambda},$$

$$e^\alpha(v \otimes e^{r+\lambda}) = v \otimes e^{\alpha+r+\lambda},$$

$$\epsilon_\alpha(v \otimes e^{r+\lambda}) = \epsilon(\alpha, r) v \otimes e^{r+\lambda},$$

$$E^\pm(\alpha, z) = \exp \left( \mp \sum_{n=1}^{\infty} \frac{z^{\mp 2n}}{n} \alpha(\pm n) \right),$$

$$F^\pm(p(\alpha), z) = \exp \left( \mp \sum_{n=1}^{\infty} \frac{2z^{\mp(2n-1)}}{2n-1} p(\alpha) \left( \pm \left( n - \frac{1}{2} \right) \right) \right),$$

and

$$\alpha_i(z) = \sum_{n \in \mathbb{Z}} \alpha_i(n) z^{-2n} + \sum_{n \in \mathbb{Z}} p(\alpha_i) \left( n - \frac{1}{2} \right) z^{-2n+1}, \quad i = 1, \dots, l,$$

$$\left( \sum_{i=1}^l c_i \alpha_i \right) (z) = \sum_{i=1}^l c_i \alpha_i(z).$$

Then  $E^\pm(\alpha, z)$  and  $F^\pm(\alpha, z)$ ,  $\alpha(z)$  are elements in  $\mathbf{End}(V(\mathcal{Q}))[[z, z^{-1}]]$ .

Let  $\widetilde{V(\mathcal{Q})}$  be the formal completion of  $V(\mathcal{Q})$ . We give vertex operators on  $\widetilde{V(\mathcal{Q})}$ :

1. For  $\alpha \in \mathcal{Q}$ , define

$$(21) \quad Y(\alpha, z) = \begin{cases} E^-(\alpha, z)E^+(\alpha, z), & \text{if } p(\alpha) = 0, \\ \sqrt{-1}E^-(\alpha, z)E^+(\alpha, z)F^-(p(\alpha), z)F^+(p(\alpha), z), & \text{if } p(\alpha) \in \Delta_S, \\ E^-(\alpha, z)E^+(\alpha, z)F^-(p(\alpha), z)F^+(p(\alpha), z), & \text{if } p(\alpha) \in \Delta_L. \end{cases}$$

2. For  $\alpha \in \mathcal{Q}$ , define

$$(22) \quad X(\alpha, z) = (-1)^{-p_0(\alpha)} Y(\alpha, z) z^{(\alpha, \alpha)} e^\alpha z^{2\alpha} \epsilon_\alpha.$$

Also define

$$(23) \quad X(a, b, z, w) = (-1)^{-p_0(a+b)} : Y(a, z) Y(b, w) : w^{(a+b, a+b)} e^{a+b} w^{a+b} \epsilon_{a+b},$$

where  $: \cdot :$  means the normal ordered product:

$$: a(m) b(n) : = \begin{cases} a(m) b(n), & m \leq n; \\ b(n) a(m), & m > n. \end{cases}$$

for suitable  $m, n \in \frac{1}{2}\mathbb{Z}$ .

3. Suppose that  $e_1, \dots, e_l$  and  $e'_1, \dots, e'_l$  are bases of  $H$  and  $H_M$ , respectively, such that

$$(e_i, e_j) = \delta_{ij}, \quad (e'_i, e'_j) = \delta_{ij}.$$

Define operator

$$(24) \quad d_0 = \frac{1}{2} \sum_{i=1}^l e_i(0) e_i(0) + \sum_{i=1}^l \sum_{n=1}^{\infty} e_i(-n) e_i(n) + \sum_{i=1}^l \sum_{n=1}^{\infty} e'_i \left( -n + \frac{1}{2} \right) e'_i \left( n - \frac{1}{2} \right).$$

For  $x = a_1(m_1) \cdots a_k(m_k) \otimes e^r \in V(\mathcal{Q})$ , define

$$\deg(x) = \sum_{j=1}^k m_j - \frac{1}{2}(r, r).$$

$\deg(x)$  is called the degree of  $x$ . Then

$$d_0(x) = -\deg(x).$$

**Lemma 4.1.**

$$(25) \quad [d_0, \alpha(m)] = -m\alpha(m),$$

$$(26) \quad [d_0, X(\alpha, z)] = \frac{1}{2} \frac{z \partial}{\partial z} X(\alpha, z).$$

*Proof.* It is clear that

$$\deg(\alpha(m) \cdot (v \otimes e^r)) = (m + \deg(v \otimes e^r))(\alpha(m) \cdot (v \otimes e^r)),$$

so

$$[d_0, \alpha(m)] \cdot (v \otimes e^r) = -m\alpha(m) \cdot (v \otimes e^r),$$

then (25) holds. From (25), we have

$$\begin{aligned} [d_0, X(\alpha, z)] &= \left( \sum_{n=1}^{\infty} \alpha(-n) z^{2n} + p(\alpha) \left( -n + \frac{1}{2} \right) z^{2n-1} + \alpha(0) + \frac{(\alpha, \alpha)}{2} \right) X(\alpha, z) \\ &\quad + X(\alpha, z) \left( \sum_{n=1}^{\infty} \alpha(n) z^{-2n} + p(\alpha) \left( n - \frac{1}{2} \right) z^{-2n+1} \right), \end{aligned}$$

which equals

$$\frac{1}{2} \frac{z \partial}{\partial z} X(\alpha, z).$$

□

**Lemma 4.2.** *For any  $\alpha \in \Delta_L \cup \Delta_M \cup \Delta_S$ , the Laurent series of  $X(\alpha, z)$  are denoted by*

$$X(\alpha, z) = \sum_{n \in \mathbb{Z}} X_{\frac{n}{2}}(\alpha) z^{-n}.$$

Particularly, if  $\alpha \in \Delta_L$ , we have

$$X(\alpha, z) = \sum_{n \in \mathbb{Z}} X_{n+\frac{1}{2}}(\alpha) z^{-n}.$$

*Proof.* Let  $r \in \mathcal{Q}$ . If  $\alpha \in \Delta_L$ , then  $\alpha = \pm 2(\alpha_i + \cdots \alpha_l)$  for some  $i > 0$ , and  $(\alpha, \lambda) = \pm \frac{1}{2}$ , for any  $r \in \mathcal{Q}$ ,  $(\alpha, r) \in \mathbb{Z}$ , so

$$\begin{aligned} 2[\deg(e^\alpha \cdot e^{\lambda+r}) - \deg(e^{\lambda+r})] &= (\alpha + r + \lambda, \alpha + r + \lambda) - (r + \lambda, r + \lambda) \\ &= (\alpha, \alpha) + 2(\alpha, r + \lambda) \\ &= 2 \pm 1 + 2(\alpha, r) \in 2\mathbb{Z} + 1, \end{aligned}$$

If  $\alpha \in \Delta_S$ , then  $\alpha = \pm(\alpha_i + \cdots \alpha_l)$  for some  $i > 0$ ,

$$\begin{aligned} 2[\deg(e^\alpha \cdot e^{\lambda+r}) - \deg(e^{\lambda+r})] &= (\alpha + r + \lambda, \alpha + r + \lambda) - (r + \lambda, r + \lambda) \\ &= (\alpha, \alpha) + 2(\alpha, r + \lambda) \\ &= \frac{1}{2} \pm \frac{1}{2} + 2(\alpha, r) \in \mathbb{Z}, \end{aligned}$$

If  $\alpha \in \Delta_M$ , then  $\alpha = \pm((\alpha_i + \cdots + \alpha_{j-1} + \alpha_j + \cdots \alpha_l) \pm (\alpha_j + \cdots + \alpha_l))$  for some  $l \geq j > i > 0$ , so  $(\alpha, \lambda) = 1$  or  $0$ , thus

$$\begin{aligned} 2[\deg(e^\alpha \cdot e^{\lambda+r}) - \deg(e^{\lambda+r})] &= (\alpha + r + \lambda, \alpha + r + \lambda) - (r + \lambda, r + \lambda) \\ &= (\alpha, \alpha) + 2(\alpha, r + \lambda) \\ &= 1 + 2(\alpha, r) \pm 1 \in \mathbb{Z} \\ \text{or} &= 1 + 2(\alpha, r) \in \mathbb{Z} \end{aligned}$$

Then by Equation(25) and the definition of  $E, F$  operators, we know that the lemma holds.  $\square$

**Theorem 4.3.** *The Lie algebra linearly generated by operators*

$$\left\{ X_{\frac{n}{2}}(\alpha), X_n(\alpha'), e_i(n), e'_i\left(n + \frac{1}{2}\right), \text{id}, d_0 \mid \alpha \in \Delta_M \cup \Delta_S, \alpha' \in \Delta_L, n \in \mathbb{Z}, i = 1, \dots, l \right\}$$

on  $V(\mathcal{Q})$  is isomorphic to the twisted affine Lie algebra of type  $A_{2l}^{(2)}$ . The isomorphism  $\pi$  is given by

$$\begin{aligned} \pi(e_\alpha \otimes s^{\frac{n}{2}}) &= X_{\frac{n}{2}}(\alpha), \quad \alpha \in \Delta_M \cup \Delta_S, \\ \pi(e_\alpha \otimes s^{n+\frac{1}{2}}) &= X_{n+\frac{1}{2}}(\alpha), \quad \alpha \in \Delta_L, \\ \pi(\gamma^{-1}(\alpha_i) \otimes s^n) &= \alpha_i(n), \\ \pi(c) &= \text{id}, \\ \pi(d) &= -d_0. \end{aligned}$$

Additionally, the image of imaginary root vectors with non-integer degree can be obtained from the above definition, together with Lie bracket.

## 5. PROOF OF THEOREM 4.3

**Lemma 5.1.**

$$[a(n), X_{\frac{m}{2}}(\alpha)] = (a, \alpha) X_{n+\frac{m}{2}}(\alpha), \quad m, n \in \mathbb{Z}.$$

By analogy of the argument in Section 3.4 of [FLM], we easily obtain by a direct calculation

**Lemma 5.2.** *Suppose that  $z$  and  $w$  are two complex variables. Then*

$$E^+(a, z)E^-(b, w) = z^{-2(a,b)}(z^2 - w^2)^{(a,b)} E^-(b, w)E^+(a, z),$$

$$F^+(p(a), z)F^-(p(b), w) = (z - w)^{(p(a), p(b))} (z + w)^{-(p(a), p(b))} F^-(p(b), w)F^+(p(a), z),$$

for  $|z| > |w|$  and  $a, b \in \Delta_L \cup \Delta_M \cup \Delta_S \cup \{0\}$ .

For complex variables  $z, w$ , in this paper,  $C_1$  means the field such that  $|z| > |w|$  and  $C_2$  means the field such that  $|z| < |w|$ .

**Lemma 5.3.** *For  $\alpha \in \Delta_S$ ,*

$$[X_{\frac{m}{2}}(\alpha), X_{\frac{n}{2}}(-\alpha)] = -2\epsilon(\alpha, -\alpha)(\delta_{m+n,0}m + 2\alpha(z)).$$

*Proof.*

$$\begin{aligned} & [X_{\frac{m}{2}}(\alpha), X(-\alpha, w)] \\ &= -\frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, -a) z^{m-1} \frac{z(z+w)}{(z-w)^2} X(a, -a, z, w) (zw^{-1})^{2a-\frac{1}{2}} dz \\ &= -2\epsilon(a, -a) w^m (m + 2\alpha(w)), \end{aligned}$$

so it is true.  $\square$

**Lemma 5.4.** *For  $\alpha \in \Delta_M$ ,*

$$[X_{\frac{m}{2}}(\alpha), X_{\frac{n}{2}}(-\alpha)] = \epsilon(\alpha, -\alpha)(\delta_{m+n,0}m + 2\alpha(z)).$$

*Proof.*

$$\begin{aligned} & [X_{\frac{m}{2}}(\alpha), X(-\alpha, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, -a) z^{m-1} \frac{z^2}{(z-w)^2} X(a, -a, z, w) (zw^{-1})^{2a-1} dz \\ &= \epsilon(a, -a) w^m (m + 2\alpha(w)), \end{aligned}$$

that is, the lemma holds.  $\square$

**Lemma 5.5.** *For  $\alpha \in \Delta_L$ ,*

$$[X_{\frac{m}{2}}(\alpha), X_{\frac{n}{2}}(-\alpha)] = \frac{1}{2}\epsilon(\alpha, -\alpha)(\delta_{m+n,0}m + 2\alpha(z)).$$

*Proof.*

$$\begin{aligned} & [X_{\frac{m}{2}}(\alpha), X(-\alpha, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, -a) z^{m-1} \frac{z^4}{(z-w)^2(z+w)^2} X(a, -a, z, w) (zw^{-1})^{2a-1} dz \\ &= \epsilon(a, -a) w^m (m + 2\alpha(w)), \end{aligned}$$

hence, the lemma holds.  $\square$

**Lemma 5.6.** *If  $a, b \in \Delta_S$  and  $a + b \in \Delta_M$ , then*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = -2\epsilon(a, b)X_{\frac{m+n}{2}}(a + b),$$

*when  $p_0(a + b) = a + b$  and*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = 2\sqrt{-1}\epsilon(a, b)X_{\frac{m+n}{2}}(a + b),$$

*when  $p_0(a + b) = a + b$ .*

*Proof.* (A) if  $p_0(a + b) = a + b$ , then  $(a, b) = 0, (p(a), p(b)) = -1$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= -\frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b)z^{m-1} \frac{z+w}{z-w} X(a, b, z, w)(zw^{-1})^{2a+\frac{1}{2}} dz \\ &= -2\epsilon(a, b)w^m X(a + b, w). \end{aligned}$$

(B) if not, we can assume that  $p_0(a + b) = b - a$ , then  $(p(a), p(b)) = 1$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= -\frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b)z^{m-1}(-1)^{-2a} \frac{z-w}{z+w} X(a, b, z, w)(zw^{-1})^{2a+\frac{1}{2}} dz \\ &= 2\sqrt{-1}\epsilon(a, b)w^m X(a + b, w), \end{aligned}$$

hence, this lemma holds.  $\square$

**Lemma 5.7.** *If  $a, b \in \Delta_S$  and  $a + b \in \Delta_L$ , then  $a = b$  and*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(a)] = (-1)^m 4\sqrt{-1}\epsilon(a, a)X_{\frac{m+n}{2}}(2a).$$

*Proof.* Now

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(a, w)] \\ &= -\frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b)z^{m-1}(-1)^{2a} \frac{(z-w)^2}{z(z+w)} X(a, b, z, w)(zw^{-1})^{2a+\frac{3}{2}} dz \\ &= (-1)^m 4\sqrt{-1}\epsilon(a, b)w^m X(2a, w), \end{aligned}$$

so this lemma is true. Note that the coefficient of bracket is not zero if and only  $m + n \in 2\mathbb{Z} + 1$ , this coincides with that

$$X(2a, z) = \sum_{n \in \mathbb{Z}} X_{n+\frac{1}{2}}(2a)z^{-2n-1}.$$

$\square$

**Lemma 5.8.** *If  $b \in \Delta_M$  and  $a, a + b \in \Delta_S$ , then*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = \epsilon(a, b)X_{\frac{m+n}{2}}(a + b).$$

*Proof.* In this case,  $p_0(a + b) = p_0(a) + p_0(b)$  and  $(a, b) = (p(a), p(b)) = -\frac{1}{2}$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(a, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b)z^{m-1} \frac{z}{(z-w)} X(a, b, z, w)(zw^{-1})^{2a-\frac{1}{2}} dz \\ &= \epsilon(a, b)w^m X(a + b, w), \end{aligned}$$

hence, the lemma is true.  $\square$



**Lemma 5.9.** *If  $a, b, a + b \in \Delta_M$ , then*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = (-1)^{2m} X_{\frac{m+n}{2}}(a + b).$$

*Proof.* (A) if  $p_0(a + b) = p_0(a) + p_0(b)$ , then  $(a, b) = (p(a), p(b)) = -\frac{1}{2}$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b) z^{m-1} \frac{z}{z-w} X(a, b, z, w) (zw^{-1})^{2a} dz \\ &= \epsilon(a, b) w^m X(a + b, w). \end{aligned}$$

(B) if not, we can assume that  $p_0(a + b) = p_0(b) - p_0(a)$ , then  $(a, b) = -\frac{1}{2}$  and  $(p(a), p(b)) = \frac{1}{2}$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b) (-1)^{-2p_0(a)} z^{m-1} (-1)^{-2a} \frac{z-w}{z+w} X(a, b, z, w) (zw^{-1})^{2a} dz \\ &= (-1)^{-2p_0(a)+2a} \epsilon(a, b) w^m X(a + b, w), \end{aligned}$$

it is clear that  $2(a - p_0(a)) \in 4\dot{\Delta} \cup \{0\}$ , so  $(-1)^{-2p_0(a)+2a} = \text{id}_{V(\mathcal{Q})}$ , hence, this lemma holds.  $\square$

**Lemma 5.10.** *If  $a, b \in \Delta_M$  and  $a + b \in \Delta_L$ , then*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = 2\epsilon(a, b) (-1)^m X_{\frac{m+n}{2}}(a + b),$$

*if  $(p(a), p(b)) = 1$  and*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = 2\epsilon(a, b) X_{\frac{m+n}{2}}(a + b),$$

*if  $(p(a), p(b)) = -1$ .*

*Proof.* In this case, there must be  $p_0(a) = \pm p_0(b)$ , (A) if  $p_0(a) = p_0(b)$ , then  $(a, b) = 0$  and  $(p(a), p(b)) = 1$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b) z^{m-1} (-1)^{-2p_0(a)} \frac{z-w}{z+w} X(a, b, z, w) (zw^{-1})^{2a} dz \\ &= 2(-1)^m \epsilon(a, b) w^m X(a + b, w) (-1)^{2(a-p_0(a))}. \end{aligned}$$

(B) if  $p_0(a) = -p_0(b)$ , then  $(a, b) = 0$  and  $(p(a), p(b)) = -1$ , so

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b) z^{m-1} \frac{z+w}{z-w} X(a, b, z, w) (zw^{-1})^{2a} dz \\ &= 2\epsilon(a, b) w^m X(a + b, w). \end{aligned}$$

$\square$

**Lemma 5.11.** *If  $b, a + b \in \Delta_M$  and  $a \in \Delta_L$ , then*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = (-1)^m X_{\frac{m+n}{2}}(a + b),$$

*when  $p_0(a) = p_0(a + b)$  and*

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = (-1)^m \frac{1 + (-1)^n}{2} X_{\frac{m+n}{2}}(a + b),$$

Here,  $m$  is odd number.

*Proof.* At first, we know that  $(a, b) = -1$ , then if  $p_0(b) = p_0(a + b)$ , we have

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(a, b) z^{m-1} \frac{z^2}{(z-w)(z+w)} X(a, b, z, w) (zw^{-1})^{2a} dz \\ &= \epsilon(a, b) w^m X(a + b, w), \end{aligned}$$

this equation hold since  $(-1)^{2a} = -\text{id}_{V(\mathcal{Q})}$ .

If  $p_0(b) = -p_0(a + b)$ , by Lemmas 5.1, 5.4 and 5.10, we have

$$\begin{aligned} & [X_{\frac{m}{2}}(a), X(b, w)] \\ &= \frac{1}{2} (-1)^m \epsilon(-b, a + b) [[X_{\frac{m}{2}}(-b), X_0(a + b)], X_{\frac{n}{2}}(b)] \\ &= \frac{1}{2} (-1)^m \epsilon(-b, a + b) \epsilon(-b, b) [w^m(m + 2b(w)), X_0(a + b)] \end{aligned}$$

since  $(b, a + b) = 0$ ,  $(p(b), p(a + b)) = -1$  and  $m$  is odd, so if  $n$  is even, we have

$$[X_{\frac{m}{2}}(a), X_{\frac{n}{2}}(b)] = (-1)^{2m} X_{\frac{m+n}{2}}(a + b),$$

otherwise, it is zero. Additionally,  $\epsilon(a, b) = 1$  since  $a \in 2\mathcal{Q}$ . Thus we have obtained the result.  $\square$

**Lemma 5.12.** *If  $b, a + b \in \Delta_S$  and  $a \in \Delta_L$ , then  $a + 2b = 0$ ,*

$$[X_{\frac{m}{2}}(-2b), X_{\frac{n}{2}}(b)] = (-1)^m \sqrt{-1} X_{\frac{m+n}{2}}(-b),$$

*Proof.*

$$\begin{aligned} & [X_{\frac{m}{2}}(-2b), X(b, w)] \\ &= \frac{1}{2\pi\sqrt{-1}} \int_{C_1-C_2} \epsilon(-2b, b) z^{m-1} \frac{z^2}{(z-w)(z+w)} X(a, b, z, w) (zw^{-1})^{-4b} dz \\ &= \sqrt{-1} \epsilon(-2b, b) w^m X(a + b, -w), \end{aligned}$$

so this lemma holds since  $\epsilon(-2b, b) = 1$ .  $\square$

By all these lemmas and Lemma 4.1, we know that Theorem 4.3 is true.

## 6. THE STRUCTURE OF $V(\mathcal{Q})$

Let

$$\alpha_0^\vee = c - \sum_{i=1}^{l-1} \alpha_i^\vee - \frac{1}{2} \alpha_l^\vee \in \mathcal{H}^\sigma.$$

Choose  $\alpha_0 \in \mathcal{H}^{\sigma*}$  such that  $\{\alpha_0, \alpha_1, \dots, \alpha_l\}$  is the simple root system of twisted affine Lie algebra  $\mathcal{G}^\sigma$  and

$$\alpha_0(d) = 1, \quad \alpha_0(\alpha_0^\vee) = 2, \quad \alpha_0(c) = 0$$

and

$$\alpha_0(\alpha_i^\vee) = -\delta_{i,1}$$

for  $i = 1, \dots, l$ . Then  $\delta = \alpha_0 + 2(\alpha_1 + \dots + \alpha_l)$  is an imaginary root of  $\mathcal{G}^\sigma$ . Let  $\Lambda_i \in \mathcal{H}^{\sigma*}$  be such that  $\Lambda_i(\alpha_j^\vee) = \delta_{i,j}$  for  $i = 0, 1, \dots, l$ .

**Lemma 6.1.**  $V(\mathcal{Q})$  is a completely reducible module and associated with Cartan subalgebra  $\mathcal{H}^\sigma$ , it has weight space decomposition

$$V(\mathcal{Q}) = \sum_{\mu \in P(V(\mathcal{Q}))} V(\mathcal{Q})_\mu.$$

The proof is very similar to those in [1],[2] and [4].

**Lemma 6.2.** If  $x$  is a highest weight vector, then it must have the form  $1 \otimes e^{\lambda+\alpha}$ .

*Proof.* It is clear since  $x$  must be commutative with  $\alpha_i(n)$  and  $p(\alpha_i)(n + \frac{1}{2})$  for any  $n \in \mathbb{Z}$  and  $i = 1, \dots, l$ .  $\square$

**Lemma 6.3.**  $1 \otimes e^\lambda$  is a highest weight vector.

*Proof.* Obviously, for any  $0 < i < l$ , we have

$$X_0(\alpha_i) \cdot (1 \otimes e^\lambda) = Y_{\frac{1}{2}}(\alpha_i) \otimes e^{\lambda+\alpha_i} = 0,$$

and

$$X_0(\alpha_l) \cdot (1 \otimes e^\lambda) = Y_{\frac{1}{2}}(\alpha_l) \otimes e^{\lambda+\alpha_l} = 0.$$

Finally,

$$X_1(-2\alpha_1 - \dots - 2\alpha_l) \cdot (1 \otimes e^\lambda) = Y_1(-2\alpha_1 - \dots - 2\alpha_l) \otimes e^{\lambda-2\alpha_1-\dots-2\alpha_l} = 0.$$

So  $1 \otimes e^\lambda$  is a highest weight vector. Particularly, by a direct computation, we know that the irreducible submodule with highest weight vector  $1 \otimes e^\lambda$  has highest weight  $\Lambda_l$ .  $\square$

**Theorem 6.4.**  $V(\mathcal{Q})$  is an irreducible  $\mathcal{G}^\sigma$ -module isomorphic to  $L(\Lambda_l)$ .

*Proof.* If  $x = 1 \otimes e^{\lambda+\alpha}$  is a highest weight vector, then for  $i = 1, \dots, l$ , by

$$X_0(\alpha_i) \cdot (x) = 0,$$

we have

$$(\alpha, \alpha_i) > -\frac{1}{2},$$

that is to say,

$$(\alpha, \alpha_i) \geq 0.$$

Secondly, by

$$X_{\frac{1}{2}}(-2\alpha_1 - \dots - 2\alpha_l) \cdot (x) = 0,$$

we have

$$(\alpha, \alpha_1 + \dots + \alpha_l) \leq \frac{1}{4}.$$

Then  $\alpha = 0$ . Thus we have proved that  $1 \otimes e^\lambda$  is the unique highest weight vector (up to a scalar). That's,  $V(\mathcal{Q})$  is irreducible.  $\square$

## REFERENCES

- [1] C. Jiang; D.J. Meng, *Vertex representations of the  $\nu + 1$ -toroidal Lie algebra of type  $B_l^{(1)}$* , J. Algebra. **246** (2001), 564–593.
- [2] D. Liu; N. Hu, *Vertex representations for toroidal Lie algebra of type  $G_2$* , Journal of Pure and Applied Algebra, **198** (2005), 257–279.
- [3] J. Lepowsky; R.Wilson, *The structure of standard modules, I: Universal algebras and the Rogers-Ramanujan identities*, Invent. math. **77** (1984), 199–290.
- [4] L. Xia; N. Hu, *Irreducible representations for Virasoro-toroidal Lie algebras*, Journal of Pure and Applied Algebra, **194** (2004), 213–237.
- [6] H. You; C. Jiang, *Vertex operators construction for the toroidal Lie algebra of type  $F_4$* , Comm. Algebra **31** (2003), 2161–2182.
- [7] V. G. Kac, *Infinite dimensional Lie algebras*, 3rd edition, Cambridge Uni. Press, Cambridge, U.K., 1990.

LX: FACULTY OF SCIENCE, JIANGSU UNIVERSITY, ZHENJIANG 212013, JIANGSU, CHINA

NH: DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200241, CHINA

XB: SCHOOL OF MATHEMATICAL SCIENCE, NANKAI UNIVERSITY, TIANJIN 300071, CHINA  
*E-mail address:* `nhhu@math.ecnu.edu.cn`